

University of Groningen

## Onderzoekings oor die werksame stowwe van verskillende gramineeënpollens volgens die Dale metode

Vermeulen, B.S.

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

1930

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Vermeulen, B. S. (1930). *Onderzoekings oor die werksame stowwe van verskillende gramineeënpollens volgens die Dale metode*. [, University of Groningen]. [S.n.].

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

## STELLINGEN.

---

### I.

N-stof bepaling van pollenextracten als maatstaf om de sterkte van de werkzame stoffen af te meten is foutief.

### II.

Roggepollenextract bevat een histamine-achtige stof.

### III.

De uitspraak van COCA en GROVE, dat de werkzame stoffen in pollens geen eiwitten zijn, is onjuist.

### IV.

Bij recidiveerende angina bij volwassenen passe men bestraling toe vóór tonsillectomie.

### V.

Het ontstaan van mediane neusfistels moet verklaard worden door afsnoering van ectodermale bestanddeelen in de sulcus supra nasalis.

### VI.

De gelijktijdige behandeling met neosalversan en bismuth volgens SHAMBERG in gevallen waarbij patient slechts eenmaal per week antiluetisch behandeld kan worden, is de aangewezen.

---

## VII.

Bezoek bij roodvonkpatienten in de barakken kan onder zekere voorwaarden worden toegestaan.

## VIII.

Baarmoedergeruisch ontstaat door torsie van de uterus.

## IX.

Behoudens enkele uitzonderingsgevallen behandel men fracturen van het onderbeen met circulair gipsverband na repositie en *wel zoo spoedig mogelijk* na het trauma.

## X.

Indien na een schedeltrauma neurologische of psychische afwijkingen gevonden worden, waarvan de aetiologie niet duidelijk is, verrichte men encephalographie.

---